arXiv:1301.4385v1 [math.CA] 18 Jan 2013

SOME INEQUALITIES FOR COMPLETE ELLIPTIC INTEGRALS

LI YIN AND FENG QI

Abstract. In the paper, by using Lupaş integral inequality, the authors find some new inequalities for the complete elliptic integrals of the first and second kinds. These results improve some known inequalities.

1. Introduction

Legendre's complete elliptic integrals of the first and second kind are defined for real numbers $0<r<1$ by

$$\kappa(r)=\int_0^{\pi/2}\frac{1}{\sqrt{1-r^2\sin^2t}}\,\mathrm{d}t=\int_0^1\frac{1}{\sqrt{(1-t^2)(1-r^2t^2)}}\,\mathrm{d}t \tag{1.1}$$

and

$$\varepsilon(r)=\int_0^{\pi/2}\sqrt{1-r^2\sin^2t}\,\mathrm{d}t=\int_0^1\sqrt{\frac{1-r^2t^2}{1-t^2}}\,\mathrm{d}t \tag{1.2}$$

respectively. They can also defined by

$$\kappa(r,s)=\int_0^{\pi/2}\frac{1}{\sqrt{r^2\cos^2t+s^2\sin^2t}}\,\mathrm{d}t \tag{1.3}$$

and

$$\varepsilon(r,s)=\int_0^{\pi/2}\sqrt{r^2\cos^2t+s^2\sin^2t}\,\mathrm{d}t. \tag{1.4}$$

Letting $r'=\sqrt{1-r^2}$. We often denote

$$\kappa'(r)=\kappa(r')\quad\text{and}\quad\varepsilon'(r)=\varepsilon(r').$$

These integrals are special cases of Gauss hypergeometric function

$$F(a,b;c;x)=\sum_{n=0}^{\infty}\frac{(a,n)(b,n)}{(c,n)}\frac{x^n}{n!},$$

where $(a,n)=\prod_{k=0}^{n-1}(a+k)$. Indeed, we have

$$\kappa(r)=\frac{\pi}{2}F\biggl(\frac12,\frac12;1;r^2\biggr)\quad\text{and}\quad\varepsilon(r)=\frac{\pi}{2}F\biggl(-\frac12,\frac12;1;r^2\biggr).$$

For more information on the history, background, properties, and applications, please refer to [4, 5] and related references.

Recently, some bounds for $\varepsilon(r)$ and $\kappa(r)$ were discovered in the paper [8]. For example, Theorem 1 in [8] states that, for $0<r<1$,

$$\frac{\pi}{2}-\frac12\ln\frac{(1+r)^{1-r}}{(1-r)^{1+r}}<\varepsilon(r)<\frac{\pi-1}{2}+\frac{1-r^2}{4r}\ln\frac{1+r}{1-r}. \tag{1.5}$$

For more information on inequalities of complete elliptic integrals, please refer to [1, 2, 3, 9] and a short survey in [10, pp. 40–46].

2010 *Mathematics Subject Classification.* Primary 33E05; Secondary 26D15, 33C75.

Key words and phrases. inequality; complete elliptic integral; Lupaş inequality.

The first author was supported partially by the Natural Science Foundation of Shandong Province under grant number ZR2012AQ028.

Motivating by the double inequality (1.5), some estimates for $\varepsilon(r)$ in terms of rational functions of the arithmetic, geometric, and roots square means were obtained in [7, 11, 12].

The aim of this paper is to establish some new inequalities for the complete elliptic integrals.

2. Lemmas

In order to prove our main results, the following lemma is necessary.

Lemma 2.1 (Lupaş integral inequality [6, p. 57])**.** *If $f', g' \in L_2[a,b]$, then*

$$\left|\frac{1}{b-a}\int_a^b f(t)g(t)\,\mathrm{d}t-\frac{1}{b-a}\int_a^b f(t)\,\mathrm{d}t\frac{1}{b-a}\int_a^b g(t)\,\mathrm{d}t\right| \le \frac{b-a}{\pi^2}\|f'\|_2\|g'\|_2, \quad (2.1)$$

where

$$\|f'\|_2=\left(\int_a^b |f'(t)|^2\,\mathrm{d}t\right)^{1/2} \quad \text{and} \quad \|g'\|_2=\left(\int_a^b |g'(t)|^2\,\mathrm{d}t\right)^{1/2}.$$

3. Main Results

Now we are in a position to find some inequalities for complete elliptic integrals.

Theorem 3.1. *For $r \in (0,1)$, we have*

$$\frac{\pi\sqrt{6+2\sqrt{1-r^2}\,-3r^2}}{4\sqrt{2}} \le \varepsilon(r) \le \frac{\pi\sqrt{10-2\sqrt{1-r^2}-5r^2}}{4\sqrt{2}}. \quad (3.1)$$

Proof. Taking

$$f(t)=g(t)=\sqrt{1-r^2\sin^2 t}$$

and letting $a=0$ and $b=\frac{\pi}{2}$ in Lemma 2.1 yield

$$\begin{aligned}\left|\frac{2}{\pi}\int_0^{\pi/2}\left(1-r^2\sin^2 t\right)\mathrm{d}t-\frac{4}{\pi^2}\varepsilon^2(r)\right| &= \left|\frac{4}{\pi^2}\varepsilon^2(r)-\frac{2-r^2}{2}\right| \\ \le \frac{1}{2\pi}\int_0^{\pi/2}\frac{r^4\sin^2 t\cos^2 t}{1-r^2\sin^2 t}\,\mathrm{d}t &= \frac{1}{2\pi}\int_0^{\pi/2} r^4\sin^2 t\cos^2 t\sum_{n=0}^{\infty} r^{2n}\sin^{2n} t\,\mathrm{d}t \\ &= \frac{1}{2\pi}\sum_{n=0}^{\infty}\int_0^{\pi/2} r^{2n+4}\left(\sin^{2n+2} t-\sin^{2n+4} t\right)\mathrm{d}t=\frac{1}{4}h(r), \quad (3.2)\end{aligned}$$

where, by

$$\int_0^{\pi/2}\sin^{2i} t\,\mathrm{d}t=\frac{\pi}{2}\frac{(2i-1)!!}{(2i)!!} \quad (3.3)$$

for $i \in \mathbb{N}$,

$$h(r)=\sum_{n=0}^{\infty}\frac{(2n+1)!!}{(2n+2)!!}\frac{r^{2n+4}}{2n+4}.$$

A direct calculation yields

$$h'(r)=\sum_{n=0}^{\infty}\frac{(2n+1)!!}{(2n+2)!!}r^{2n+3}=r\sum_{n=0}^{\infty}\frac{(2n+1)!!}{(2n+2)!!}r^{2n+2}=r\left(\frac{1}{\sqrt{1-r^2}}-1\right),$$

where we used

$$\frac{1}{\sqrt{1-t^2}}=\sum_{n=0}^{\infty}\frac{(2i-1)!!}{(2i)!!}t^{2n}, \quad |t|<1. \quad (3.4)$$

Hence, we have

$$h(r) = h(0) + \int_0^r h'(r)\,\mathrm{d}r = 1 - \sqrt{1-r^2} - \frac{r^2}{2}. \tag{3.5}$$

Substituting this equality into (3.2) gives

$$\left|\frac{4}{\pi^2}\varepsilon^2(r) - \frac{2-r^2}{2}\right| \le \frac{1}{4} - \frac{\sqrt{1-r^2}}{4} - \frac{r^2}{8}.$$

This means the double inequality (3.1). □

Remark 3.1. By the well-known software Mathematica, we can show that

1. the left-hand side inequality in (3.1) refines the corresponding one in (1.5);
2. the right-hand side inequalities in (3.1) and (1.5) are not contained each other;
3. when $r \in [\frac{1}{4}, \frac{3}{4}]$, the right-hand side inequality in (3.1) is better than the corresponding one in (1.5).

Theorem 3.2. *For $r \in (0,1)$, we have*

$$\frac{\pi\sqrt{32 - r^4 - 32r^2}}{8\sqrt{2}\,\sqrt[4]{(1-r^2)^3}} \le \kappa(r) \le \frac{\pi\sqrt{r^4 - 32r^2 + 32}}{8\sqrt{2}\,\sqrt[4]{(1-r^2)^3}}. \tag{3.6}$$

Proof. Taking

$$f(t) = g(t) = \frac{1}{\sqrt{1 - r^2\sin^2 t}}$$

and letting $a = 0$ and $b = \frac{\pi}{2}$ in Lemma 2.1 lead to

$$\begin{aligned}
\left|\frac{2}{\pi}\int_0^{\pi/2}\frac{1}{1-r^2\sin^2 t}\,\mathrm{d}t - \frac{4}{\pi^2}\kappa^2(r)\right| &= \left|\frac{4}{\pi^2}\kappa^2(r) - \frac{2}{\pi}\int_0^{\pi/2}\sum_{n=0}^{\infty} r^{2n}\sin^{2n} t\,\mathrm{d}t\right| \\
&= \left|\frac{4}{\pi^2}\kappa^2(r) - \sum_{n=0}^{\infty}\frac{(2n-1)!!}{(2n)!!}r^{2n}\right| = \left|\frac{4}{\pi^2}\kappa^2(r) - \frac{1}{\sqrt{1-r^2}}\right| \\
&\le \frac{1}{2\pi}\int_0^{\pi/2}\frac{r^4\sin^2 t\cos^2 t}{\left(1-r^2\sin^2 t\right)^3}\,\mathrm{d}t \\
&= \frac{1}{4\pi}\int_0^{\pi/2} r^4\sin^2 t\cos^2 t\sum_{n=0}^{\infty}(n+2)(n+1)r^{2n}\sin^{2n} t\,\mathrm{d}t \\
&= \frac{1}{8}\sum_{n=0}^{\infty}(n+2)(n+1)r^{2n+4}\frac{(2n+1)!!}{(2n+2)!!}\frac{1}{2n+4} = \frac{r^3}{32}p(r),
\end{aligned}$$

where

$$p(r) = \sum_{n=0}^{\infty}\frac{(2n+2)(2n+1)!!}{(2n+2)!!}r^{2n+1}$$

satisfies $p(0) = 0$,

$$\int_0^r p(r)\,\mathrm{d}r = \sum_{n=0}^{\infty}\frac{(2n+1)!!}{(2n+2)!!}r^{2n+2} = \frac{1}{\sqrt{1-r^2}} - 1,$$

and so

$$p(r) = \frac{r}{\sqrt{(1-r^2)^3}}. \tag{3.7}$$

Consequently, we find

$$\left|\frac{4}{\pi^2}\kappa^2(r) - \frac{1}{\sqrt{1-r^2}}\right| \le \frac{r^4}{32\sqrt{(1-r^2)^3}}. \tag{3.8}$$

The double inequality (3.6) follows. ☐

Theorem 3.3. *For $r > 0$ and $s > 0$, we have*

$$\frac{\pi}{8}\sqrt{\frac{8rs(r^2+s^2)-(s^2-r^2)^2}{rs}} \le \varepsilon(r,s) \le \frac{\pi}{8}\sqrt{\frac{8rs(r^2+s^2)+(s^2-r^2)^2}{rs}}\,. \tag{3.9}$$

Proof. Taking

$$f(t)=g(t)=\sqrt{r^2\cos^2 t+s^2\sin^2 t}$$

and letting $a=0$ and $b=\frac{\pi}{2}$ in Lemma 2.1 reveal

$$\begin{aligned}\biggl|\frac{2}{\pi}\int_0^{\pi/2}\bigl(r^2\cos^2 t+s^2\sin^2 t\bigr)\,\mathrm{d}\,t-\frac{4}{\pi^2}\varepsilon^2(r,s)\biggr| &= \biggl|\frac{4}{\pi^2}\varepsilon^2(r,s)-\frac{r^2+s^2}{2}\biggr| \\ \le \frac{1}{2\pi}\int_0^{\pi/2}\frac{(s^2-r^2)^2\sin^2 t\cos^2 t}{r^2\cos^2 t+s^2\sin^2 t}\,\mathrm{d}\,t &\le \frac{(s^2-r^2)^2}{2\pi}\frac{1}{4}\int_0^{\pi/2}\frac{1}{r^2\cos^2 t+s^2\sin^2 t}\,\mathrm{d}\,t \\ = \frac{(s^2-r^2)^2}{8\pi}\frac{1}{rs}\arctan\biggl(\frac{s}{r}\tan t\big|_0^{\pi/2}\biggr) &= \frac{(s^2-r^2)^2}{16rs}.\end{aligned}$$

This means the double inequality (3.9). ☐

Theorem 3.4. *For $s > r > 0$, we have*

$$\biggl|\frac{4}{\pi^2}\kappa^2(r,s)-\frac{1}{\pi rs}\biggr| \le \frac{s^2-r^2}{32rs}\biggl(\frac{1}{s^2}+\frac{1}{r^2}\biggr). \tag{3.10}$$

Proof. Taking

$$f(t)=g(t)=\frac{1}{\sqrt{r^2\cos^2 t+s^2\sin^2 t}}$$

and letting $a=0$ and $b=\frac{\pi}{2}$ in Lemma 2.1 figure out

$$\begin{aligned}\biggl|\frac{2}{\pi}\int_0^{\pi/2}\frac{1}{r^2\cos^2 t+s^2\sin^2 t}\,\mathrm{d}\,t-\frac{4}{\pi^2}\kappa^2(r,s)\biggr| &= \biggl|\frac{4}{\pi^2}\kappa^2(r,s)-\frac{1}{\pi rs}\biggr| \\ &\le \frac{1}{2\pi}\int_0^{\pi/2}\frac{\bigl(s^2-r^2\bigr)^2\sin^2 t\cos^2 t}{\bigl(r^2\cos^2 t+s^2\sin^2 t\bigr)^3}\,\mathrm{d}\,t \\ &= \frac{r^2-s^2}{8\pi}\int_0^{\pi/2}\sin t\cos t\,\mathrm{d}\,\frac{1}{\bigl(r^2\cos^2 t+s^2\sin^2 t\bigr)^2} \\ &= \frac{r^2-s^2}{8\pi}\biggl[\int_0^{\pi/2}\frac{\sin^2 t}{\bigl(r^2\cos^2 t+s^2\sin^2 t\bigr)^2}\,\mathrm{d}\,t-\int_0^{\pi/2}\frac{\cos^2 t}{\bigl(r^2\cos^2 t+s^2\sin^2 t\bigr)^2}\,\mathrm{d}\,t\biggr] \\ &= \frac{r^2-s^2}{8\pi}\biggl[\int_0^{\pi/2}\frac{\csc^2 t}{\bigl(r^2\cot^2 t+s^2\bigr)^2}\,\mathrm{d}\,t-\int_0^{\pi/2}\frac{\sec^2 t}{\bigl(r^2+s^2\tan^2 t\bigr)^2}\,\mathrm{d}\,t\biggr] \\ &= \frac{r^2-s^2}{8\pi}\biggl[\int_0^{\infty}\frac{1}{(r^2u^2+s^2)^2}\,\mathrm{d}\,u-\int_0^{\pi/2}\frac{1}{(r^2+s^2\lambda^2)^2}\,\mathrm{d}\,\lambda\biggr]=\frac{s^2-r^2}{32rs}\biggl(\frac{1}{s^2}+\frac{1}{r^2}\biggr).\end{aligned}$$

The proof is complete. ☐

Remark 3.2. From (3.10), we easily obtain

$$\frac{\pi}{2}\sqrt{\frac{32r^2s^2-\pi(s^4-r^4)}{32\pi r^3s^3}} \le \kappa(r,s) \le \frac{\pi}{2}\sqrt{\frac{32r^2s^2+\pi(s^4-r^4)}{32\pi r^3s^3}}\,. \tag{3.11}$$

By the software MATHEMATICA, we can show that the double inequality in (3.11) and the inequality in [8, Theorem 2] are not contained each other.

References

[1] H. Alzer, *Sharp inequalities for the complete elliptic integrals of the first kinds*, Math. Proc. Camb. Phil. Soc. **124** (1988), no. 2, 309–314; Available online at `http://dx.doi.org/10.1017/S0305004198002692`. 1

[2] G. D. Anderson, P. Duren, and M. K. Vamanamurthy, *An inequality for complete elliptic integrals*, J. Math. Anal. Appl. **182** (1994), no. 1, 257–259; Available online at `http://dx.doi.org/10.1006/jmaa.1994.1080`. 1

[3] G. D. Anderson, S.-L. Qiu, and M. K. Vamanamurthy, *Elliptic integrals inequalities, with applications*, Constr. Approx. **14** (1998), no. 2, 195–207; Available online at `http://dx.doi.org/10.1007/s003659900070`. 1

[4] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen, *Topics in special functions*, available online at `http://arxiv.org/abs/0712.3856`. 1

[5] G. D. Anderson, M. Vuorinen, and X.-H. Zhang, *Topics in special functions III*, available online at `http://arxiv.org/abs/1209.1696`. 1

[6] N. S. Barnett and S. S. Dragomir, *On an inequality of the Lupaş type*, Demonstratio Math. **42** (2008), no. 1, 57–62. 2

[7] Y.-M. Chu, M.-K. Wang, S.-L. Qiu, and Y.-P. Jiang, *Bounds for complete elliptic integrals of second kind with applications*, Comput. Math. Appl. **63** (2012), 1177–1184; Available online at `http://dx.doi.org/10.1016/j.camwa.2011.12.038`. 2

[8] B.-N. Guo and F. Qi, *Some bounds for the complete elliptic integrals of the first and second kind*, Math. Inequal. Appl. **14** (2011), no. 2, 323–334; Available online at `http://dx.doi.org/10.7153/mia-14-26`. 1, 4

[9] F. Qi and Z. Huang, *Inequalities for complete elliptic integrals*, Tamkang J. Math. **29** (1998), no. 3, 165–169; Available online at `http://dx.doi.org/10.5556/j.tkjm.29.1998.165-169`. 1

[10] F. Qi, D.-W. Niu, and B.-N. Guo, *Refinements, generalizations, and applications of Jordan's inequality and related problems*, J. Inequal. Appl. **2009** (2009), Article ID 271923, 52 pages; Available online at `http://dx.doi.org/10.1155/2009/271923`. 1

[11] M.-K. Wang and Y.-M. Chu, *Asymptotical bounds for complete elliptic integrals of second kind*, available online at `http://arxiv.org/abs/1209.0066`. 2

[12] M.-K. Wang, Y.-M. Chu , S.-L. Qiu, and Y.-P. Jiang, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164** (2012), 928–937; Available online at `http://dx.doi.org/10.1016/j.jat.2012.03.011`. 2

(Yin) Department of Mathematics, Binzhou University, Binzhou City, Shandong Province, 256603, China
E-mail address: `yinli_79@163.com`

(Qi) School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China
E-mail address: `qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com`
URL: `http://qifeng618.wordpress.com`